

## Leucistic Marbled Salamanders (*Ambystoma opacum*) in Virginia

Joseph C. Mitchell

Department of Biology  
University of Richmond  
Richmond, Virginia 23173

Don R. Church

Department of Biology  
University of Virginia  
Charlottesville, Virginia 22903

Albinism and its various forms have been reported for a variety of species and life history stages of North American amphibians (Hensley, 1959; Brame, 1962; Dyrkacz, 1981; Bechtel, 1995). Most instances of albinism are based on single individuals in populations that otherwise contain normally pigmented individuals. Examples from Virginia include American Toad (*Bufo americanus*) from Fairfax County, Upland Chorus Frog (*Pseudacris feriarum*) from Frederick County, Eastern Spadefoot (*Scaphiopus holbrookii*) from New Kent County, Northern Gray-cheeked Salamander (*Plethodon montanus*; originally reported as *P. jordani*) from Smyth County, and Peaks of Otter Salamander (*Plethodon hubrichti*) from Botetourt County (Hensley, 1959; Bulmer, 1975; Hayslett et al., 1998). In some cases, however, numerous albinistic individuals are reported for a single population. An increase in frequency of albinism in the Shovel-nosed Salamander (*Desmognathus marmoratus*) was reported for a population in North Carolina studied over a six-year period by Martof (1962) and Martof & Walton (1965). Albinism or its more common leucistic form (lacking integumentary pigment with normal eyes; Dyrkacz, 1981) has been reported for the five species of amphibians in Virginia noted above. None of the reports, however, suggest more than single occurrences within populations.

Albinism is known to occur in 8 of the 14 recognized species of salamanders in the genus *Ambystoma* (exclusive of the unisexual forms) throughout North America (Dyrkacz, 1981; Petranka, 1998). In the mid-Atlantic region, reports of albinistic ambystomatids include larval Jefferson Salamanders (*A. jeffersonianum*), an adult Spotted Salamander (*A.*

*maculatum*), and larval Marbled Salamanders (*A. opacum*) in Maryland (Harris, 1967, 1968, 1970), and larval Mole Salamanders (*A. talpoideum*) from North Carolina (Palmer & Braswell, 1980). Dodd (1977-78) illustrated a leucistic Marbled Salamander larva but provided no locality data. No occurrences of albinism have been reported for mole salamanders in Virginia. The purpose of this note is to report two instances of the leucistic phase of albinism in *Ambystoma opacum* from the upper Coastal Plain of Virginia and the Shenandoah Valley.

On 22 April 1998, a leucistic *Ambystoma opacum* larva was collected with a dip net in a small, vegetated, vernal pool (formerly a road rut) on Fort A.P. Hill Military Reservation, Caroline County, Virginia (Fig. 1). It completely lacked dark pigment on the body and tail but had normally colored (dark) eyes. The gills were pink. Measurements of this larva were 27 mm snout-vent length (SVL), 49 mm total length, and 0.86 g. A sample of normally pigmented larvae caught in the same pool on the same day averaged ( $\pm$  1 SD) 30.6  $\pm$  1.5 mm SVL (27-35, n = 36), 55.0  $\pm$  3.3 mm total length (50-61, n = 31), and 1.3  $\pm$  0.2 g (0.98-1.64, n = 36). The leucistic larva was smaller than the rest of the cohort in all respects. It was kept alive for photographs but subsequently lost. Other aquatic amphibians found on the same date in the pool were Spotted Salamander larvae, Red-spotted Newt (*Notophthalmus viridescens*) adults, and Green Frog (*Rana clamitans*) tadpoles; all had normal pigmentation.

On 14 June 2002, a leucistic *Ambystoma opacum* metamorph was captured in a pitfall trap leaving Pond 2 in the Maple Flats Pond Complex within the Shenandoah Valley Sinkhole Pond system of the Big

Levels area in the George Washington National Forest of southeastern Augusta County, Virginia (see Buhlmann et al., 1999). This animal lacked dark pigment on the body and had only a narrow line of dark pigment on the dorsal ridge of the tail (Fig. 2). The eyes and gill stubs were dark. The leucistic metamorph measured 39 mm SVL and 69 mm TL and weighed 1.56 g. Twelve normal metamorphs captured in the same pitfall on this date averaged  $42.6 \pm 1.2$  mm SVL (38-45),  $71.9 \pm 2.1$  mm TL (62-76), and  $1.71 \pm 0.2$  g (1.56-2.94). Other amphibians found in Pond 2 during June 2002 included Spring Peeper (*Pseudacris crucifer*) larvae and a few adult Red-spotted Newts. No other ambystomatids bred in Pond 2 in the 2001-02 season. The leucistic salamander was released.

We examined over 4,279 larvae (789 in Caroline County, 3,490 in the Shenandoah Valley) and 12,167 metamorphs (Shenandoah Valley) of *Ambystoma opacum* from these two locations during 1997-2002. All but these two individuals exhibited the normal pigmented phenotype. Genetically-based albinism and other pigment abnormalities occur naturally but rarely in most vertebrates, including humans. Our observations and literature review suggest that the leucistic form of albinism occurs rarely in *A. opacum* populations. The two reported here provide the first documentation of leucistic marbled salamanders in Virginia.

#### ACKNOWLEDGMENTS

We thank C. Todd Georgel for his assistance in the field at Fort A.P. Hill, and especially Heather Mansfield and Terry Banks of the Environment and Natural Resources section of Fort A. P. Hill's Department of Public Works for support and funding of the field work in which the larvae were discovered. Funding was administered by Jeff Walden and his staff at the Conservation Management Institute (formerly the Fish and Wildlife Information Exchange), Virginia Polytechnic Institute and State University. Funding for Church's study in the Shenandoah Valley was provided by the U.S. Forest Service. USFS biologists Fred Huber, Mike Donahue, and Dawn Kirk provided helpful logistical support.

#### LITERATURE CITED

- Bechtel, H. B. 1995. Reptile and Amphibian Variants, Colors, Patterns, and Scales. Krieger Publishing Co., Malabar, FL. 206 pp.
- Brame, A. H., Jr. 1962. A survey of albinism in salamanders. *Abhandlungen und Berichte Naturkunde Vorgesichte* 11: 65-81.
- Bulmer, W. 1975. Albino toad from No. VA. Virginia Herpetological Society Bulletin 78: 7.
- Buhlmann, K. A., J. C. Mitchell, & L. R. Smith. 1999. Descriptive ecology of the Shenandoah Valley Sinkhole Pond system in Virginia. *Banisteria* 13: 23-51.
- Dodd, C. K., Jr. 1977-78. Amphibians and reptiles - the declining species. *Water Spectrum* 10: 24-32.
- Dyrkacz, S. 1981. Recent instances of albinism in North American amphibians and reptiles. Society for the Study of Amphibians and Reptiles, Herpetological Circular 11: 1-31.
- Harris, H. S., Jr. 1967. Albinism in the Jefferson Salamander, *Ambystoma jeffersonianum* Green. *Bulletin of the Maryland Herpetological Society* 3: 99-100.
- Harris, H. S., Jr. 1968. A survey of albinism in Maryland populations of amphibians and reptiles. *Bulletin of the Maryland Herpetological Society* 4: 57-60.
- Harris, H. S., Jr. 1970. Abnormal pigmentation in Maryland amphibians and reptiles. *Bulletin of the Maryland Herpetological Society* 6: 21-27.
- Hayslett, M. S., G. W. Wilson, & J. C. Mitchell. 1998. Natural history: *Plethodon hubrichti*. Albinism. *Herpetological Review* 29: 229-230.
- Hensley, M. 1959. Albinism in North American amphibians and reptiles. *Publications of the Museum, Michigan State University* 1: 133-159.
- Martof, B. S. 1962. Some aspects of the life history and ecology of the salamander *Leurognathus marmoratus*. *American Midland Naturalist* 67: 1-35.
- Martof, B. S., & K. G. Walton. 1965. A study of the unusual pigmentation in the salamander *Leurognathus marmoratus*. *Physiological Zoology* 38: 349-352.
- Palmer, W. M., & A. L. Braswell. 1980. Additional records of albinistic amphibians and reptiles from North Carolina. *Brimleyana* 3: 49-52.
- Petranka, J. W. 1998. Salamanders of the United States and Canada. Smithsonian Institution Press, Washington, DC. 587 pp.



Fig. 1. Leucistic and normal *Ambystoma opacum* larvae from Fort A.P. Hill, Caroline County, Virginia.




Fig. 2. Leucistic and normal *Ambystoma opacum* metamorphs from the Shenandoah Valley sinkhole ponds, Augusta County, Virginia.