

SHORTER CONTRIBUTIONS

Banisteria, Number 29, pages 32-35
© 2007 Virginia Natural History Society

MACROLOPHUS BREVICORNIS ON *TRIOSTEUM PERFOLIATUM* IN VIRGINIA: STATE RECORD AND NEW HOST-PLANT RECORD FOR MIRIDAE (HEMIPTERA: HETEROPTERA) -- Many dicyphine plant bugs—mirids of the subfamily Bryocorinae, tribe Dicyphini, subtribe Dicyphina (*sensu* Schuh, 1995)—are omnivores on glandular-hairy (“sticky”) plants. The bugs not only feed on their hosts but also prey on small arthropods and scavenge insects that alight on the plants and become entrapped in glandular exudations (Spomer, 1999; Wheeler, 2001). Several dicyphines native to Europe have been used in greenhouses to control pests such as aphids, thrips, and whiteflies. The Palearctic bugs used in biological control are prohibited from being introduced into North America; as omnivores, they could become plant pests (e.g., Alomar & Albajes, 1996; Wheeler, 2001; Sanchez et al., 2004). North American entomologists, therefore, have investigated native dicyphines as potential biocontrol agents (McGregor et al., 1999, 2000; Sanchez et al., 2003, 2004; McGregor & Gillespie, 2004).

Recently, the Nearctic dicyphine *Dicyphus vestitus* Uhler was recorded from heart-leaved skullcap, *Scutellaria ovata* J. Hill (Lamiaceae), in Virginia shale barrens. Virginia represented the southeasternmost record of this mirid in the eastern United States, and heart-leaved skullcap was the first host plant documented for this bug (Henry, 1999).

On 7 May 1993, I discovered another little-known dicyphine that proved to be a new record for Virginia when Thomas Rawinski took me to a dolomite glade (“Dixie Cliff”) near Glenvar in Roanoke County. I was interested in the glade because of its colonies of moss phlox, *Phlox subulata* L., a mat-forming member of the Polemoniaceae that harbors a diverse insect fauna (Wheeler, 1995a, b). During the visit to Dixie Cliff, I found late instars of a dicyphine mirid in thin woods at the base of the glade’s south-facing slope (37° 15.2' N, 80° 10.4' W). Nymphs were on horse-gentian (also known as feverwort or wild coffee), *Triosteum perfoliatum* L. (Caprifoliaceae). I was not aware of mirids known from this plant and wanted to determine the bug’s identity.

Fifth instars were collected and held at 22-24°C in a small plastic box (ca. 8 x 2 cm) with excised horse-gentian shoots; the stems were wrapped with moist cotton. Adults appeared on 13 May. I returned to Dixie Cliff in late May to collect additional adults, which Thomas Henry identified as *Macrolophus brevicornis*

Knight. Voucher specimens have been deposited in the National Museum of Natural History, Smithsonian Institution, Washington, DC.

Macrolophus Fieber is a Holarctic genus that includes five Nearctic species. The species in eastern North America, in addition to *M. brevicornis*, are *M. separatus* Uhler and *M. tenuicornis* Blatchley (Henry & Wheeler, 1988; Schuh, 1995; Maw et al., 2000). Knight (1926) provided a key to the three eastern species.

Macrolophus brevicornis was described by Knight (1926) from Iowa, Kansas, Maryland, Missouri, and New Jersey. Since the original description, only Illinois (Knight, 1941) and Kentucky (Henry et al., 2005) have been added to the distribution. Biological information on this seldom-collected mirid is limited to its collection from two plant species: a milkweed, *Asclepias* sp. (Asclepiadaceae), in Iowa (Knight, 1941), and a ground-cherry, *Physalis* sp. (Solanaceae), in Kentucky (Henry et al., 2005). Knight (1941) noted “breeding” on milkweed, implying that nymphs were observed. Certain true bugs, or heteropterans, such as lygaeine Lygaeidae (Price & Willson, 1979; Slater & Baranowski, 1990), specialize on asclepiads, but specialization by North American mirids on these cardenolide-rich plants is unknown (Knight, 1941, 1968; Schuh, 1995; Wheeler, 2001).

Additional observations are needed to determine if *M. brevicornis* actually develops on milkweeds; the bug apparently has not been recorded from *Asclepias* since Knight’s (1941) report. *Macrolophus brevicornis*, however, has been collected consistently on glandular species of *Physalis* (A.G.W., unpubl. data). *Triosteum perfoliatum* apparently can be added to the bug’s known host-plant range. I also found two adults on this plant in a Maryland shale barren (Washington Co., Sideling Hill Wildlife Management Area, near Little Orleans) in mid-June 1994.

The mirid’s seasonal history on *T. perfoliatum* is incompletely known. Third through fifth instars were observed at Dixie Cliff in early May 1993; by 28 May, adults predominated (23 were collected) with fewer than 10 fifth instars found. No individuals of *M. brevicornis* were observed on 3 July. In 1994, fourth and fifth instars were found on 20 May, mostly on lower (abaxial) leaf surfaces. Whether more than one generation is produced is unknown. *Macrolophus tenuicornis*, the only eastern species of the genus that has been studied, is bivoltine in Pennsylvania (Wheeler et al., 1979).

Horse-gentian is a densely glandular-hairy herb (Gleason & Cronquist, 1991; Rhoads & Block, 2000), although its exudate appears not to entrap small

arthropods. Glandular trichomes presumably evolved as a defense against herbivory by small arthropods (Levin, 1973; Duffey, 1986; Gregory et al., 1986; Sugiura & Yamazaki, 2006) and provide greater protection than nonglandular trichomes (Duke, 1994; van Dam & Hare, 1998). Glandular trichomes are toxic to certain herbivores and can deter oviposition, impede movement, and alter feeding behavior, as well as the searching behavior of their natural enemies. Sticky plants tend to be “off limits” to most generalist herbivores, which cannot traverse the glandular surfaces. Benefits to plants from reduced herbivory can be offset by the adverse effects of glandular trichomes on herbivores’ parasitoids and predators (van Dam & Hare, 1998; Gassmann & Hare, 2005). Plants might also benefit by digesting and absorbing proteins from trapped insects (protocarnivory) (Spomer, 1999) or obtaining supplemental nitrogen when trapped insects decay and the breakdown products leach into the soil (Eisner, 2003).

Omnivorous insects generally have broader host-plant ranges than strict herbivores (Eubanks et al., 2003). Dicyphine mirids, as well as many stilt bugs (Berytidae) (Wheeler & Schaefer, 1982; Henry, 1997; Henry & Froeschner, 1998), associated with glandular-pubescent plants tend to be omnivores and might be more polyphagous than members of their respective families that develop on nonglandular hosts. The berytid *Jalysus spinosus* (Say), for example, feeds on unrelated sticky dicots and monocots (Wheeler & Henry, 1981; Wheeler, 1986, 1994). Studies on omnivorous heteropterans that specialize on sticky plants, similar to the study of terrestrial heteropterans that are omnivores (Eubanks et al., 2003), likely would yield insights into the ecological significance of associations with glandular plants and the evolutionary consequences of such specialization.

ACKNOWLEDGMENTS

I thank Thomas J. Rawinski (formerly with Division of Natural Heritage, Virginia Department of Conservation and Recreation, Richmond) for taking me to Dixie Cliff and identifying *Triosteum perfoliatum*, Thomas J. Henry (Systematic Entomology Laboratory, USDA, ARS, PSI, c/o National Museum of Natural History, Smithsonian Institution, Washington, DC) for identifying *Macrolophus brevicornis* and reviewing the manuscript, and Peter H. Adler (Department of Entomology, Soils, and Plant Sciences, Clemson University) for his review of an earlier draft of the manuscript.

LITERATURE CITED

- Alomar, O., & R. Albajes. 1996. Greenhouse whitefly (Homoptera: Aleyrodidae) predation and tomato fruit injury by the zoophytophagous predator *Dicyphus tamaninii* (Heteroptera: Miridae). Pp. 155-177 *In* O. Alomar & R. N. Wiedenmann (eds.), *Zoophytophagous Heteroptera: Implications for Life History and Integrated Pest Management*. Thomas Say Publications in Entomology: Proceedings. Entomological Society of America, Lanham, MD.
- Duffey, S. S. 1986. Plant glandular trichomes: their partial role in defence against insects. Pp. 151-172 *In* B. Juniper & T. R. E. Southwood (eds.), *Insects and the Plant Surface*. Edward Arnold, London.
- Duke, S. O. 1994. Glandular trichomes—a focal point of chemical and structural interactions. *International Journal of Plant Sciences* 155: 617-620.
- Eisner, T. 2003. *For Love of Insects*. Harvard University Press, Cambridge, MA. 448 pp.
- Eubanks, M. D., J. D. Styrsky, & R. F. Denno. 2003. The evolution of omnivory in heteropteran insects. *Ecology* 84: 2549-2556.
- Gassmann, A. J., & J. D. Hare. 2005. Indirect cost of a defensive trait: variation in trichome type affects the natural enemies of herbivorous insects on *Datura wrightii*. *Oecologia* 144: 62-71.
- Gleason, H. A., & A. Cronquist. 1991. *Manual of Vascular Plants of Northeastern United States and Adjacent Canada*. Second edition. New York Botanical Garden, Bronx, NY. 910 pp.
- Gregory, P., D. A. Avé, P. Y. Bouthyette, & W. M. Tingey. 1986. Insect-defensive chemistry of potato glandular trichomes. Pp. 173-183 *In* B. Juniper & T. R. E. Southwood (eds.), *Insects and the Plant Surface*. Edward Arnold, London.
- Henry, T. J. 1997. Monograph of the stilt bugs, or Berytidae (Heteroptera), of the Western Hemisphere. *Memoirs of the Entomological Society of Washington* 19: 1-149.
- Henry, T. J. 1999. Review of the eastern North American *Dicyphus*, with a key to species and redescription and neotype designation for *D. vestitus*

- Uhler (Heteroptera: Miridae). Proceedings of the Entomological Society of Washington 101: 832-838.
- Henry, T. J., & R. C. Froeschner. 1998. Catalog of the stilt bugs, or Berytidae, of the world (Insecta: Hemiptera: Heteroptera). Contributions of the American Entomological Institute 30(4): 1-72.
- Henry, T. J., & A. G. Wheeler, Jr. 1988. Family Miridae Hahn, 1833 (=Capsidae Burmeister, 1835). The plant bugs. Pp. 251-507 In T. J. Henry & R. C. Froeschner (eds.), Catalog of the Heteroptera, or True Bugs, of Canada and the Continental United States. E. J. Brill, Leiden.
- Henry, T. J., C. V. Covell, Jr., & A. G. Wheeler, Jr. 2005. An annotated list of the plant bugs, or Miridae (Hemiptera: Heteroptera), of Kentucky. Journal of the New York Entomological Society 113: 24-76.
- Knight, H. H. 1926. A key to the North American species of *Macropsophus* with descriptions of two new species (Hem.: Miridae). Entomological News 37: 313-316.
- Knight, H. H. 1941. The plant bugs, or Miridae, of Illinois. Illinois Natural History Survey Bulletin 22(4): 1-234.
- Knight, H. H. 1968. Taxonomic review: Miridae of the Nevada Test Site and the western United States. Brigham Young University Science Bulletin. Biological Series 9(3): 1-282.
- Levin, D. A. 1973. The role of trichomes in plant defense. Quarterly Review of Biology 48: 3-15.
- Maw, H. E. L., R. G. Foottit, K. G. A. Hamilton, & G. G. E. Scudder. 2000. Checklist of the Hemiptera of Canada and Alaska. NRC Research Press, Ottawa, ON. 220 pp.
- McGregor, R. R., & D. R. Gillespie. 2004. Olfactory responses of the omnivorous generalist predator *Dicyphus hesperus* to plant and prey odours. Entomologia Experimentalis et Applicata 112: 201-205.
- McGregor, R. R., D. R. Gillespie, D. M. J. Quiring, & M. R. J. Foisy. 1999. Potential use of *Dicyphus hesperus* Knight (Heteroptera: Miridae) for biological control of pests of greenhouse tomatoes. Biological Control 16: 104-110.
- McGregor, R. R., D. R. Gillespie, C. G. Park, D. M. J. Quiring, & M. R. J. Foisy. 2000. Leaves or fruit? The potential for damage to tomato fruits by the omnivorous predator, *Dicyphus hesperus*. Entomologia Experimentalis et Applicata 95: 325-328.
- Price, P. W., & M. F. Willson. 1979. Abundance of herbivores on six milkweed species in Illinois. American Midland Naturalist 101: 76-86.
- Rhoads, A. F., & T. A. Block. 2000. The Plants of Pennsylvania: an Illustrated Manual. University of Pennsylvania Press, Philadelphia. 1,061 pp.
- Sanchez, J. A., D. R. Gillespie, & R. R. McGregor. 2003. The effects of mullein (*Verbascum thapsus*) on the population dynamics of *Dicyphus hesperus* (Heteroptera: Miridae) in tomato greenhouses. Biological Control 28: 313-319.
- Sanchez, J. A., D. R. Gillespie, & R. R. McGregor. 2004. Plant preference in relation to life history traits in the zoophytophagous predator *Dicyphus hesperus*. Entomologia Experimentalis et Applicata 112: 7-19.
- Schuh, R. T. 1995. Plant Bugs of the World (Insecta: Heteroptera: Miridae): Systematic Catalog, Distributions, Host List, and Bibliography. New York Entomological Society, New York. 1,329 pp.
- Slater, J. A., & R. M. Baranowski. 1990. Lygaeidae of Florida (Hemiptera: Heteroptera). Arthropods of Florida and Neighboring Land Areas. Vol. 14. Florida Department of Agriculture and Consumer Services, Gainesville. 211 pp.
- Spomer, G. G. 1999. Evidence of protocarnivorous capabilities in *Geranium viscosissimum* and *Potentilla arguta* and other sticky plants. International Journal of Plant Sciences 160: 98-101.
- Sugiura, S. & K. Yamazaki. 2006. Consequences of scavenging behaviour in a plant bug associated with a glandular plant. Biological Journal of the Linnean Society 88: 593-602.
- van Dam, N. M., & J. D. Hare. 1998. Differences in distribution and performance of two sap-sucking herbivores on glandular and non-glandular *Datura wrightii*. Ecological Entomology 23: 22-32.

Wheeler, A. G., Jr. 1986. A new host association for the stilt bug *Jalysus spinosus* (Heteroptera: Berytidae). *Entomological News* 97: 63-65.

Wheeler, A. G., Jr. 1994. A new host for *Jalysus spinosus* (Heteroptera: Berytidae) and new host family (Commelinaceae) for stilt bugs. *Entomological News* 105: 201-203.

Wheeler, A. G., Jr. 1995a. Plant bugs (Heteroptera: Miridae) of *Phlox subulata* and other narrow-leaved phloxes in eastern United States. *Proceedings of the Entomological Society of Washington* 97: 435-451.

Wheeler, A. G., Jr. 1995b. Insects of moss phlox (*Phlox subulata*): unexpected diversity in Appalachian shale barrens. *Virginia Journal of Science* 46: 148 (Abstract).

Wheeler, A. G., Jr. 2001. *Biology of the Plant Bugs (Hemiptera: Miridae): Pests, Predators, Opportunists*. Cornell University Press, Ithaca, NY. 507 pp.

Wheeler, A. G., Jr., & T. J. Henry. 1981. *Jalysus spinosus* and *J. wickhami* (Hemiptera: Berytidae): taxonomic clarification, review of host plants and distribution, and keys to adults and fifth instars. *Annals of the Entomological Society of America* 74: 606-615.

Wheeler, A. G., Jr., & C. W. Schaefer. 1982. Review of stilt bug (Hemiptera: Berytidae) host plants. *Annals of the Entomological Society of America* 75: 498-506.

Wheeler, A. G., Jr., G. L. Miller, & T. J. Henry. 1979. Biology and habits of *Macrolophus tenuicornis* (Hemiptera: Miridae) on hayscentedfern (Pteridophyta: Polypodiaceae). *Melsheimer Entomological Series* 27: 11-17.

Alfred G. Wheeler, Jr.
Department of Entomology, Soils, and Plant Sciences
Clemson University
Clemson, South Carolina 29634-0315